THE SIDEREAL ZODIAC, EXPLAINED

The Complete Guide to Sidereal Ayanamsa



History, competing systems, the 2026 sky, and the choice that makes the most sense

Agastya Trivedi

Vedic Astrologer

2026 EDITION

A note from the author

If you have ever changed the ayanamsa setting in an astrology app and watched the chart rearrange itself, you already know why this subject can become strangely emotional. One option keeps the Moon in Dhanishta. Another moves it into Shatabhisha. One gives a planet a comfortable Navamsha. Another places it across the boundary. Both calculations may be astronomically competent, yet they cannot both be using the same zodiac origin.

This guide is my attempt to make the debate understandable without flattening it. New students will find the basic astronomy, practical examples, and a clear choice. Experienced astrologers will find the historical variants, exact 2026 values, boundary timing, software naming problems, and the reasons why phrases such as True Lahiri can be misleading.

Ayanamsa is not a contest between people who respect tradition and people who respect astronomy. The real task is to name the rule, calculate it consistently, understand what it changes, and avoid claiming more evidence than we actually possess.

> The short answer
> If you need one default in 2026, use standard Lahiri, specifically the IAE 1985 implementation. It is the most practical common language for Vedic astrology. If you deliberately want the rule that keeps Spica exactly at 0 degrees Libra on the date of calculation, use True Chitra and label it clearly. No system has been proven to be the universal predictive winner.

Agastya Trivedi

Vedic Astrologer

Contents

A note from the author 2

1. The one setting that quietly changes the whole chart 4

2. The astronomy, without the unnecessary fog 4

3. The Lahiri family: familiar, useful, and less simple than it looks 5

4. True Chitra and the problem with the label True Lahiri 6

5. KP ayanamsa: why a few minutes become a big deal 7

6. Raman, Pushya, and the search for a better anchor 8

7. Mula, Revati, Dhruva, and the galactic imagination 8

8. Surya Siddhanta, Aryabhata, Yukteswar, and historical zero years 10

9. Fagan-Bradley and the Babylonian line 10

10. The 2026 comparison: the same sky through different grids 11

11. Ingress timing in 2026: hours become days, and days become months 13

12. Where the differences matter most in a natal chart 14

13. How to test an ayanamsa without fooling yourself 15

14. A decision guide for astrologers and software builders 16

15. Questions astrologers ask after changing the setting 17

16. The conclusion: which ayanamsa makes the most sense in 2026? 19

Appendix A. Extended 2026 values 21

Appendix B. Glossary 21

START HERE

1. The one setting that quietly changes the whole chart

Ayanamsa is the angular distance between the tropical zodiac and a chosen sidereal zodiac. Tropical 0 degrees Aries follows the moving March equinox. Sidereal 0 degrees Aries is tied to a stellar, historical, or geometric reference. In 2026 the familiar Lahiri offset is a little over 24 degrees. The practical conversion is simple: subtract the chosen ayanamsa from the tropical longitude and wrap the answer back into the 360 degree circle.

> A quick example
> At the research snapshot, the tropical Sun was at 4 degrees 30 minutes Virgo. Subtract the Lahiri ayanamsa of about 24 degrees 14 minutes and the same physical Sun becomes approximately 10 degrees 17 minutes Leo in the sidereal zodiac. Nothing in the sky moved. Only the zero point changed.

That sounds like bookkeeping until a point is close to a boundary. Then the consequences multiply. The Moon can change nakshatra and alter the starting Vimshottari dasha. A planet can change pada and therefore Navamsha. The Ascendant can move into another sign. A slow-planet ingress forecast can shift by days or even months. In KP, where sub-lords subdivide the zodiac very finely, a few arcminutes may affect the result even when the sign stays the same.

There are also two separate questions that are often mixed together. First, can astronomy calculate the planet and the chosen reference accurately? Usually yes. Second, which reference should define the sidereal zodiac? Astronomy cannot answer that on its own because Spica, Revati, Pushya, Mula, a historical equinox, and the Galactic Center are different proposed anchors.

SOURCE 1: Swiss Ephemeris sidereal documentation
The most useful technical map of modern ayanamsa implementations. It explains the formulas, epochs, star anchors, software modes, proper motion, and known naming ambiguities.
https://www.astro.com/ftp/swisseph/doc/swisseph.htm

A useful mental model is to imagine a transparent zodiac wheel laid over the sky. Every ayanamsa system agrees on the positions of the physical bodies. Each system rotates the transparent wheel to a different starting line. The debate is about where that line belongs.

THE MOVING REFERENCE FRAME

2. The astronomy, without the unnecessary fog

Precession is the slow change in the orientation of Earth's rotation axis. It causes the equinox to move westward relative to the stellar background. People often reduce this to a fixed 25,800

year cycle or a neat annual rate, but modern astronomy uses time-dependent models. The equator changes, the ecliptic changes, and the motion is not a perfectly rigid clock.

This matters because historical ayanamsas often begin with a zero year and then add a constant number of arcseconds per year. Such formulas can work as approximations near the intended era, but they are not identical to a modern precession model. A software label can therefore conceal a larger question: is it preserving the founder's table, the founder's approximate rule, a later reconstruction, or a modern astronomical anchor?

SOURCE 2: IAU 2006 Resolution B1
The International Astronomical Union adopted the P03 precession theory and gave a precise modern definition of the ecliptic. This is why contemporary astronomy does not rely on one eternal annual precession rate.
https://www.iers.org/fileadmin/SharedDocs/Publikationen/EN/IERS/resolutions/IAU/IAU2006_Resolution_B1.pdf

Nutation adds a smaller, shorter-period oscillation on top of precession. That is why software may offer a mean ayanamsa and an apparent or true value. At the 2026 snapshot used in this guide, adding nutation altered the comparable value by roughly 9.6 arcseconds. That is tiny for a sign, but it is not tiny if a program claims arcsecond precision.

Stars also have proper motion. A star is called fixed only in the everyday astrological sense. If your definition says Spica must remain exactly at 0 degrees Libra, the clean modern method is to calculate Spica's tropical longitude for the date and derive the ayanamsa from it. An old epoch plus a constant rate will slowly drift away from that literal promise.

Projection is another hidden choice. A star away from the ecliptic can be projected to ecliptic longitude in more than one way. Galactic systems are especially sensitive to this. The ordinary ecliptic longitude of the Galactic Center, a polar projection through the celestial pole, and the intersection of the galactic plane with the ecliptic are not interchangeable.

Do not confuse signs with constellations
The twelve astrological signs are equal 30 degree sectors. The modern IAU constellations are unequal regions with official boundaries. A planet can be in a sidereal sign while visually appearing against a differently named constellation. A telescope image cannot settle the ayanamsa question by itself.

INDIA'S MODERN STANDARD

3. The Lahiri family: familiar, useful, and less simple than it looks

Most people encounter Lahiri as though it were one timeless number discovered by one astrologer. The history is more interesting. The Chitra idea, placing Spica at or near the beginning of sidereal Libra, predates N. C. Lahiri. Modern standardization came through India's Calendar Reform Committee, chaired by Meghnad Saha, with Lahiri serving as a member and becoming the name most closely associated with the result.

SOURCE 3: Report of the Calendar Reform Committee, 1955
The government report documents the effort to unify India's divergent calendar practices. It is the institutional background for the modern standard that became widely known as Lahiri or Chitrapaksha.
https://archive.org/details/dli.csl.963

The committee's goal was not to run an astrology tournament. It needed a consistent framework for calendars, tithis, nakshatras, festivals, and ephemerides. That distinction is important. Standardization gives us continuity and interoperability. It does not automatically prove that the chosen zero point predicts every event better than all rivals.

The standard modern implementation used in major software is commonly identified as Lahiri IAE 1985. At 2026-08-27 18:39:49 UTC its mean ayanamsa was 24 degrees 13 minutes 45.97 seconds. Nearby variants include the pre-1985 ICRC value, Lahiri 1940, and a reconstructed zero-year-285 model. They differ by seconds or about a minute, which is normally invisible in sign-level work but can matter near fine boundaries.

SOURCE 4: Rashtriya Panchang, Government of India
The official national panchanga illustrates how a standardized nirayana framework supports calendar practice. It is evidence of institutional continuity, not a controlled test of astrological prediction.
https://mausam.imd.gov.in/responsive/rashtriyPanchang.php

SOURCE 5: Indian Astronomical Ephemeris
The Positional Astronomy Centre describes India's official astronomical ephemeris work. It helps separate high-quality planetary calculation from the separate choice of sidereal zero.
https://mausam.imd.gov.in/responsive/indianAstronomicalEphemeris.php

Why does standard Lahiri remain so powerful in practice? Because an astrologer rarely works alone. We compare charts with books, teachers, panchangas, software, research archives, and other practitioners. A shared default reduces invisible disagreements. When somebody says that the Moon is in a particular nakshatra without naming the ayanamsa, Lahiri is usually the least surprising assumption in mainstream Vedic work.

My practical reading
Lahiri is not the winner because every rival has been disproved. It is the winner as a default because it combines a clear modern implementation, enormous ecosystem support, continuity with Indian ephemeris practice, and a difference from True Chitra that is small enough to manage with boundary warnings.

LITERAL STAR ANCHORING

4. True Chitra and the problem with the label True Lahiri

True Chitra asks a beautifully direct question: what if Chitrapaksha means exactly what it says? Calculate Spica's current tropical ecliptic longitude and subtract 180 degrees so that Spica sits exactly at 0 degrees Libra in the sidereal zodiac. This dynamically handles proper motion and the changing coordinate frame.

At the 2026 snapshot, True Chitra was 24 degrees 12 minutes 34.87 seconds, only 1 minute 11.10 seconds below standard Lahiri. Most charts will look identical. The disagreement appears near a nakshatra, pada, Navamsha, cusp, or other fine boundary.

This makes True Chitra the most astronomically exact implementation of one specific rule: Spica must remain at exactly 180 degrees sidereal longitude. It does not make the rule itself scientifically proven as the unique astrological origin. Astronomical exactness is always exactness relative to a declared anchor.

The phrase True Lahiri should be treated cautiously. Depending on the software or community, it may mean True Chitra, Lahiri 1940, the year-285 reconstruction, or a compatibility setting inherited from another program. A serious chart record should name the formula or software mode, not rely on that ambiguous label.

> Safe label
> Use True Chitra, Spica fixed at 0 degrees Libra. Avoid a bare True Lahiri option. The first is reproducible. The second may change meaning from one program to another.

KRISHNAMURTI PADDHATI

5. KP ayanamsa: why a few minutes become a big deal

K. S. Krishnamurti built KP around stellar lords and unequal sub-lord divisions. This makes boundary precision operationally important. In ordinary sign-based reading, a difference of five arcminutes may do nothing. In KP it can change a sub-lord and therefore the chain of significators used for judgment and timing.

Two modern KP values deserve separate names. The table-derived KP implementation, commonly preserved as software mode 5, gave 24 degrees 07 minutes 57.31 seconds at the 2026 snapshot. The reconstructed VP291 implementation gave 24 degrees 09 minutes 09.75 seconds. Their difference is about 1 minute 12 seconds. Both are several minutes below standard Lahiri.

The reason for the duplication is historical. Krishnamurti referred to year 291 for zero ayanamsa, while his printed tables and numerical practice are not perfectly captured by a bare year. Later researchers reconstructed an exact March equinox date and modern precession treatment. The result is clearer mathematically, but it is not numerically identical to the inherited table.

SOURCE 6: Study of KP Ayanamsa with Modern Precession Theories
D. Senthilathiban reconstructs the year-291 proposal using modern precession models and explains why the table and the zero-year interpretation do not collapse into one value.
https://www.logicastro.com/files/A-Study-of-KP-Ayanamsa-D-Senthilathiban.pdf

For KP practitioners
Do not write only KP ayanamsa in exported charts. Store either KP table or KP VP291, along with the software version. If a sensitive sub-lord changes, show both values before rectifying the birth time.

IMPORTANT INDIAN ALTERNATIVES

6. Raman, Pushya, and the search for a better anchor

B. V. Raman's ayanamsa is one of the best-known Indian alternatives. It is smaller than Lahiri by about 1 degree 26 minutes 47 seconds in the 2026 snapshot. Raman presented a practical zero-year and annual-rate calculation in his books. Yet printed editions and later technical summaries do not always agree on the exact zero-year reading. Modern software resolves that uncertainty by preserving a defined computational mode.

Raman often changes padas and sometimes nakshatras without producing the dramatic rotation seen in the most distant galactic systems. In the 2026 sky snapshot, the Moon moved from Dhanishta under Lahiri to Shatabhisha under Raman, while the Sun and Mercury remained in Magha.

True Pushya Paksha was proposed by P. V. R. Narasimha Rao in 2013. It fixes Delta Cancri, Asellus Australis, at 16 degrees Cancer. The argument combines comparisons with siddhantic star positions, the place of Pushya in the nakshatra tradition, symbolism linking Cancer and the heart, and astrological case studies.

SOURCE 7: Introducing Pushya-paksha Ayanamsa
This is Rao's own proposal and therefore the best place to understand the anchor, textual comparisons, symbolic reasoning, and example charts in the form he intended.
https://www.vedicastrologer.org/articles/pp_ayanamsa.pdf

True Pushya is admirably clear as a modern definition. Independent programmers can reproduce it, and Delta Cancri's position can be recomputed rather than approximated from a zero year. Its 2026 value was 23 degrees 05 minutes 22.57 seconds, about 1 degree 08 minutes below Lahiri.

The limitation is evidence, not coherence. The textual fit is not perfect for every ancient star position, symbolic reasoning cannot uniquely choose a coordinate origin, and case studies selected after the fact do not establish universal predictive superiority. Pushya remains a serious research alternative, especially for astrologers committed to Rao's framework, but it is not yet a responsible universal default.

WHERE THE ZODIAC BEGINS

7. Mula, Revati, Dhruva, and the galactic imagination

Some systems ask a more radical question than Lahiri versus Raman. They look toward the center of the Milky Way, the root symbolism of Mula, or the final star of Revati and argue that the zodiac should be oriented around a cosmic axis rather than Spica.

True Mula, associated with K. Chandra Hari's work in the 1990s, places Lambda Scorpii at 0 degrees Sagittarius. The proposal draws on a reinterpretation of the Surya Siddhanta, the meaning of Mula as root, and arguments about the origin of the zodiac. At the 2026 snapshot it was about 43 minutes 50 seconds above Lahiri.

SOURCE 8: On the Origin of Sidereal Zodiac and Astronomy
K. Chandra Hari's history-of-astronomy paper presents the intellectual background behind the Mula-centered reconstruction. It is valuable precisely because the historical claim can be evaluated separately from later predictive enthusiasm.
https://insa.ndl.gov.in/items/f5e35867-83ed-4f9d-8693-8f35106baa58

True Revati fixes Zeta Piscium at 29 degrees 50 minutes Pisces. Its supporters point to the traditional end of Revati and to strands of Greek, Persian, Arabic, and Indian astronomical continuity. In 2026 its ayanamsa was about 3 degrees 48 minutes below Lahiri. Usha-Shashi and Sassanian variants sit in the same broad family but do not use exactly the same offset.

Dhruva Galactic Center Middle Mula, proposed by Ernst Wilhelm, uses a polar projection of the Galactic Center through the celestial north pole and places the projected point at the middle of Mula, 6 degrees 40 minutes Sagittarius. This is not the same geometry as simply taking the Galactic Center's ordinary ecliptic longitude. The 2026 Dhruva value was 20 degrees 26 minutes 10.14 seconds, nearly 3 degrees 48 minutes below Lahiri and very close to True Revati numerically.

The ordinary Galactic Center at 0 degrees Sagittarius system goes in the other direction. It had an ayanamsa of 27 degrees 13 minutes 34.19 seconds, almost 3 degrees above Lahiri. The anchor is easy to explain and compute, but it is a modern proposal. The age of the galaxy does not automatically make the astrological convention ancient.

Other galactic proposals include Rafael Gil Brand's golden-section construction, Raymond Mardyks' Galactic Alignment model, Nick Anthony Fiorenza's galactic-equator convention, Ardra Galactic Plane, and David Cochrane's Galactic Center at 0 degrees Capricorn. Each defines a different relationship between the ecliptic, the galactic plane, cardinal points, and symbolic zodiac structure.

SOURCE 9: The Cochrane Ayanamsha
David Cochrane presents a modern Galactic Center at 0 degrees Capricorn model and explicitly frames empirical testing as necessary. That honesty is useful when assessing any new ayanamsa proposal.
https://astrosoftware.com/cpnew/a_and_v/ayanamsha.html

The key caution
A precise galactic geometry can be astronomically elegant and still lack ancient Vedic provenance or predictive validation. Geometry answers how to compute the anchor. It does not prove why astrology must use it.

8. Surya Siddhanta, Aryabhata, Yukteswar, and historical zero years

A zero-year ayanamsa says that tropical and sidereal 0 degrees Aries coincided at a chosen historical date. From that date, precession is accumulated forward or backward. The attraction is obvious: find the original coincidence and the zodiac seems solved. The problem is that historical texts may use mean planets, different precession concepts, different star projections, and cycles designed for calculation rather than a modern coordinate frame.

There is no single uncontested Surya Siddhanta ayanamsa. Software modes may reconstruct an equinox near 499 CE, use a star-based interpretation, or preserve a traditional constant. Aryabhata variants face similar questions, including whether the intended coincidence belongs near 499 or 522 CE. Two programs can both say Surya Siddhanta while implementing different assumptions.

Sri Yukteswar's ayanamsa is linked to the cosmology and yuga framework in The Holy Science. It is often summarized through a Revati-oriented zero around 499 CE and an annual rate near 54 arcseconds. The text's numerical statements do not form one perfectly consistent modern formula, so software implementations depend on interpretation. Its 2026 value in a common implementation was 22 degrees 51 minutes 04.13 seconds.

SOURCE 10: The Holy Science
The authorized digital edition is the correct primary source for understanding Yukteswar's own yuga and astronomical premises. Modern software values should be described as interpretations of that framework.
https://bookstore.yogananda-srf.org/wp-content/uploads/pdf/The-Holy-Science_.pdf

Historical zero-year systems deserve study, but their age does not remove the need for precise metadata. Record the epoch date, timescale, precession model, and whether the calculation refers to a mean or true equinox. A year alone is not enough for arcminute work.

WESTERN SIDEREAL TRADITIONS

9. Fagan-Bradley and the Babylonian line

Fagan-Bradley is not a Vedic standard, but it matters because it shaped twentieth-century western sidereal astrology and appears in nearly every serious ephemeris library. Cyril Fagan and Donald Bradley argued for a Babylonian-rooted uniform zodiac and a synetic vernal point. In its common form, Spica sits at 29 degrees 06 minutes 05 seconds Virgo rather than exactly at 0 degrees Libra.

Its 2026 ayanamsa was 25 degrees 06 minutes 45.52 seconds, about 53 minutes above Lahiri. Aldebaran at exactly 15 degrees Taurus gives a very similar result, as do scholarly Babylonian reconstructions associated with Peter Huber. John P. Britton's later tablet analysis produces another nearby historical estimate.

SOURCE 11: Primer of Sidereal Astrology
Fagan and R. C. Firebrace set out the modern western sidereal framework. The book is important for intellectual history, though its astrological claims are not a validation of Vedic techniques.
https://archive.org/details/primerofsidereal0000faga

These systems remind us that a sidereal zodiac is not uniquely Indian. Ancient cultures used stellar reference frameworks in different ways. They also remind us not to blend traditions carelessly. A Babylonian historical reconstruction can be strong history without automatically becoming the best zero for Vimshottari dasha, divisional charts, or Parashari house lordship.

NUMBERS THAT MAKE THE DEBATE REAL

10. The 2026 comparison: the same sky through different grids

The following values use the exact snapshot from the supplied research: 2026-08-27 18:39:49 UTC. They are mean ayanamsas without nutation. The planetary examples use direct sidereal calculations. Values are rounded for reading, so they should not be treated as a substitute for an ephemeris API.

System	2026 mean value	From Lahiri	Anchor or use
Lahiri IAE 1985	24d 13m 45.97s	0m 00.0s	Mainstream Vedic default
True Chitra	24d 12m 34.87s	-1m 11.1s	Spica exactly at 0 Libra
KP table	24d 07m 57.31s	-5m 48.7s	Inherited KP table
KP VP291	24d 09m 09.75s	-4m 36.2s	Reconstructed zero year
Raman	22d 46m 59.29s	-1d 26m 46.7s	Raman software standard
True Pushya	23d 05m 22.57s	-1d 08m 23.4s	Delta Cancri at 16 Cancer
Dhruva GC Middle Mula	20d 26m 10.14s	-3d 47m 35.8s	Polar GC at middle Mula
GC at 0 Sagittarius	27d 13m 34.19s	+2d 59m 48.2s	Ordinary GC longitude
Fagan-Bradley	25d 06m 45.52s	+52m 59.5s	Western sidereal standard
True Revati	20d 25m 17.03s	-3d 48m 28.9s	Zeta Piscium anchor
True Mula	24d 57m 35.83s	+43m 49.9s	Lambda Scorpii at 0 Sagittarius

Ayanamsa values are circular offsets. The chart below shows the main cluster from roughly 20 to 27 degrees. Cochrane's wrapped value near 357 degrees is omitted because plotting it on the same linear scale would be misleading.



The nearest Lahiri variants differ by seconds or a few arcminutes. True Mula and Fagan-Bradley are less than a degree above it. Raman, Pushya, and Yukteswar are more than a degree below it. True Revati and Dhruva sit almost four degrees below it. The ordinary Galactic Center at 0 degrees Sagittarius sits almost three degrees above it.

SOURCE 12: Astrodienst Lahiri sidereal ephemeris for 2026
The published 2026 ephemeris provides an external reference for the contemporary Lahiri sky and is generated with Swiss Ephemeris and modern JPL planetary data.
https://www.astro.com/swisseph/slae/2000/slae_2026.pdf

What changed in the nakshatras?

Body	Lahiri	True Chitra	Pushya	Dhruva	GC 0 Sag
Moon	Dhanishta p4	Dhanishta p4	Shatabhisha p1	Shatabhisha p1	Dhanishta p3
Sun	Magha p4	Magha p4	Magha p4	Purva Phalguni p1	Magha p3
Mercury	Magha p4	Magha p4	Magha p4	Purva Phalguni p1	Magha p3
Venus	Chitra p1	Chitra p1	Chitra p1	Chitra p2	Hasta p4
Mars	Ardra p3	Ardra p3	Ardra p4	Punarvasu p1	Ardra p3
Jupiter	Ashlesha p1	Ashlesha p1	Ashlesha p1	Ashlesha p2	Pushya p4
Saturn	Revati p1	Revati p1	Revati p2	Revati p3	Revati p1
Pluto	Uttara Ashadha p4	Uttara Ashadha p4	Shravana p1	Shravana p1	Uttara Ashadha p3

At this instant all selected planets remained in the same 30 degree signs across eight representative systems. That was luck, not equivalence. The Moon, Sun, Mercury, Venus, Mars, Jupiter, Uranus, and Pluto already showed nakshatra or pada differences. A sign-only comparison would have missed the entire practical dispute.

Separate issue: mean node versus true node
The choice between mean Rahu and true Rahu can create another boundary difference. Do not blame every node discrepancy on ayanamsa. A complete chart record must state both choices.

THE SPEED OF THE PLANET MATTERS

11. Ingress timing in 2026: hours become days, and days become months

An angular difference becomes a time difference according to the body's speed. The fast Moon turns a degree into roughly two hours. Venus and the Sun turn a few degrees into days. Jupiter can turn the same spread into weeks or months, especially near a station.

Moon: Dhanishta to Shatabhisha

System	UTC crossing in 2026
Dhruva	27 Aug, 13:34
Raman	27 Aug, 18:01
True Pushya	27 Aug, 18:36
KP table	27 Aug, 20:34
True Chitra	27 Aug, 20:43
Lahiri	27 Aug, 20:45
Fagan-Bradley	27 Aug, 22:25
GC 0 Sagittarius	28 Aug, 02:24

The first and last Moon crossings were separated by about 12 hours 51 minutes. For a birth in that window, different systems can assign a different janma nakshatra, starting dasha lord, dasha balance, and panchanga label.

Venus: Virgo to Libra

System	First direct UTC crossing in 2026
Dhruva	28 Aug, 15:51
Raman	31 Aug, 12:26
True Pushya	31 Aug, 21:36
KP table	2 Sep, 05:16
True Chitra	2 Sep, 07:37
Lahiri	2 Sep, 08:14
Fagan-Bradley	3 Sep, 11:40
GC 0 Sagittarius	6 Sep, 07:47

The Venus spread was about 8 days 16 hours. A transit article that says only Venus enters Libra on 2 September is incomplete. It should also say Lahiri, True Chitra, KP, or another named ayanamsa.

Jupiter: Cancer to Leo

System	First direct crossing
Dhruva	5 Oct 2026
Raman	20 Oct 2026
True Pushya	22 Oct 2026
KP table	30 Oct 2026
True Chitra	31 Oct 2026
Lahiri	31 Oct 2026
Fagan-Bradley	8 Nov 2026
GC 0 Sagittarius	12 Jul 2027

The ordinary Galactic Center system did not produce this ingress in 2026 at all. It placed the first crossing in July 2027. This is why slow-planet transit narratives can become almost unrecognizable across distant ayanamsas.

SOURCE 13: MoonDasha Lahiri Transit Checker
This Lahiri-based transit checker is a useful example tool for seeing current sidereal positions. When comparing any tool with another source, confirm that both use the same ayanamsa, node type, location convention, and timestamp.
https://moondasha.github.io/vedic-planetary-transits/

BOUNDARY SENSITIVITY

12. Where the differences matter most in a natal chart

Ayanamsa does not change every technique equally. Angular relationships between planets remain mostly unchanged because every longitude is rotated by the same amount. Sign lordship, nakshatras, padas, and divisional-chart mappings can change sharply because they depend on boundaries.

Nakshatra and dasha balance

A nakshatra spans 13 degrees 20 minutes. A one-degree change moves a point through 7.5 percent of a nakshatra. In Vimshottari dasha, that alters the unelapsed fraction of the birth nakshatra. If the nakshatra lord owns a 20-year mahadasha, one degree corresponds to about 1.5 years of starting balance. Crossing the nakshatra boundary changes the starting lord itself, which is far more consequential.

Pada and Navamsha

Each nakshatra pada spans 3 degrees 20 minutes and maps to a Navamsha sign. A one-degree ayanamsa difference is 30 percent of a pada. Even when the D1 sign stays fixed, the D9 placement can change. This is one reason astrologers may disagree about marriage, dharma, dignity, and the maturity of a planet while apparently using the same birth data.

Higher divisional charts

Higher vargas divide a sign into increasingly narrow segments. D60 changes every 30 arcminutes. D144 changes every 12 minutes 30 seconds. At that scale, almost every major ayanamsa choice produces a different chart. These charts also demand exceptional birth-time accuracy. A precise ayanamsa cannot rescue an uncertain birth time.

Ascendant and houses

The Ascendant moves roughly one degree every four minutes on average, though the real rate varies by latitude and sign. Near a sign boundary, a modest ayanamsa change may flip the Lagna. House systems add another choice. Whole-sign houses follow the sign change directly, while cusp-based methods introduce separate geometry.

KP cusps and sub-lords

KP is the natural stress test. Sub-lord boundaries are fine enough that a few arcminutes can change the result. If the cusp or significator is close to a boundary, calculate it under the exact KP variant used by the teacher or rule set. Do not silently substitute Lahiri because the two charts look almost the same.

> Best practice
> Every serious chart engine should calculate the angular distance from each sensitive point to the nearest sign, nakshatra, pada, and requested varga boundary. If an alternative ayanamsa would cross that boundary, show a visible warning instead of pretending that the result is certain.

A PRACTICAL RESEARCH METHOD

13. How to test an ayanamsa without fooling yourself

Ayanamsa research easily turns into narrative matching. We look at a famous life, try several systems, notice the one that makes a striking yoga, and remember the success. The misses disappear. This is not dishonesty. It is normal human pattern recognition operating without guardrails.

- Write the hypothesis before opening the charts. Name the systems, techniques, outcomes, and scoring rules.
- Use accurately timed charts that were not rectified to fit the preferred theory.
- Separate a large-difference experiment, such as Lahiri versus Dhruva, from a fine-difference experiment, such as Lahiri versus True Chitra.
- Freeze the interpretive rules. Do not change yogas, aspect rules, orb rules, or event definitions after seeing the answer.
- Score real outcomes that another researcher can code, not impressions such as powerful energy or spiritual depth.

- Keep a held-out sample. A system chosen on one set of charts must predict another set it has never seen.
- Publish misses, boundary cases, and ambiguous birth times alongside successes.

The 2026 transit windows in this guide offer a clean prospective opportunity. Before the ingress, astrologers can publish dated expectations under each ayanamsa. The outcome can then be evaluated without moving the target. One transit will not settle the zodiac, but a repeated, predeclared program could gradually improve the evidence.

SOURCE 14: NASA JPL Horizons System
Horizons is an authoritative reference for high-precision solar-system ephemerides. It can verify the astronomical starting positions, but it cannot decide which astrological sidereal zero point should be preferred.
https://ssd.jpl.nasa.gov/horizons/manual.html

Scientific testing of astrology as a whole has a difficult and contested history. The lesson for ayanamsa research is modest: a confident case narrative is not the same as a blinded comparative test. We can practice astrology seriously while being precise about the kind of evidence we possess.

SOURCE 15: A double-blind test of astrology, Nature
Shawn Carlson's 1985 study is often cited in broader discussions of astrological testing. It did not compare ayanamsas, but it illustrates why blinding, preregistration, and explicit scoring matter when claims are evaluated.
https://doi.org/10.1038/318419a0

CHOOSE BY METHOD, THEN DOCUMENT IT

14. A decision guide for astrologers and software builders

Use case	Recommended setting	Reason
General Vedic natal and transit work	Lahiri IAE 1985	Best interoperability with books, teachers, panchangas, and software
Literal Spica-at-0-Libra practice	True Chitra	Most exact implementation of that declared star rule
Krishnamurti Paddhati	KP table or KP VP291	Follow the exact lineage or software standard and name the variant
P. V. R. Narasimha Rao methods	True Pushya	The author's techniques were developed around this anchor
B. V. Raman lineage	Raman	Preserves continuity with Raman's interpretive corpus
Western sidereal astrology	Fagan-Bradley	Common western sidereal reference
Experimental Mula or galactic work	Named chosen system	Use as a research model, not an unnamed correction to Lahiri

For software, the visible label should never be the only record. Store the ayanamsa identifier, exact value at the chart epoch, mean or apparent status, precession model, star catalogue or epoch if relevant, node type, ephemeris backend, coordinate frame, observer convention, timestamp, and software version.

If the user changes ayanamsa, recalculate everything downstream. Do not merely rotate the D1 planets. Nakshatra, pada, dashas, Vargas, house positions, yogas, strength calculations, and transit dates may all depend on the new longitude.

Safe names

- Lahiri, IAE 1985 standard
- True Chitra, Spica exactly at 0 degrees Libra
- KP table
- KP VP291
- Raman, software mode stated
- True Pushya, Delta Cancri at 16 degrees Cancer
- Dhruva GC Middle Mula, polar projection
- Galactic Center at 0 degrees Sagittarius, ordinary ecliptic longitude

Unsafe names include True Lahiri, Vedic ayanamsa, Galactic ayanamsa, and Surya Siddhanta without a formula. These labels sound definite while hiding multiple incompatible implementations.

A PRACTICAL FAQ

15. Questions astrologers ask after changing the setting

Does ayanamsa change planetary aspects?

The angular separation between two planets remains almost the same because the same offset is applied to both. Sign-based aspects, sign lordship, dignity, nakshatra relationships, and house placement can change because those techniques depend on the rotated zodiac grid. A geometric conjunction does not disappear merely because the sign label changes, but its dispositor and sign context may change.

Is Lahiri officially approved, and does that prove it is correct?

Lahiri has continuity with India's Calendar Reform process, national panchanga practice, and astronomical ephemerides. That makes it a powerful common standard. Government adoption solved a calendar coordination problem. It was not a blinded competition between astrological predictions, so official use should not be presented as proof of universal predictive superiority.

Is True Chitra simply a more accurate Lahiri?

It is more exact only if the rule is defined as Spica always being at 0 degrees Libra. Standard Lahiri follows a standardized ephemeris convention and differs slightly from the star's dynamically recalculated longitude. Calling one accurate and the other inaccurate hides the fact that they preserve slightly different definitions.

Can I identify the right ayanamsa by seeing which natal chart feels like me?

You can use personal resonance to generate a hypothesis, but not to finish the test. Astrological symbolism is flexible enough that several charts can sound convincing. Test dated events, freeze the rules before interpretation, and include failures. A chart that feels right may reflect a useful system, a skilled reader, or selective attention.

Should I switch ayanamsa for different techniques?

Only when the technique belongs to a school that explicitly requires its own ayanamsa. KP should use a named KP variant. A Pushya-based method should use True Pushya. For a unified Parashari practice, changing the ayanamsa each time a result looks uncomfortable creates an unfalsifiable system. Keep one default and disclose deliberate exceptions.

Why do two Lahiri apps disagree by a few seconds or minutes?

Possible causes include the Lahiri variant, mean versus apparent ayanamsa, nutation, ephemeris backend, Delta T, timescale conversion, topocentric versus geocentric coordinates, rounding, and the treatment of the lunar node. The software name alone is not enough to diagnose the difference.

Does ayanamsa affect the tropical chart?

No. The tropical zodiac is anchored to the equinox and does not need a sidereal offset. Ayanamsa is used when converting a tropical ecliptic longitude into a selected sidereal framework.

Can today's visible constellations reveal the correct sidereal zodiac?

Not directly. Astrological signs are twelve equal sectors. IAU constellations are unequal regions. Visual placement can check a star or planet's direction, but it cannot decide which equal-sector boundary astrology should adopt.

Why are divisional charts so unstable?

A high varga maps very small slices of a sign into a new zodiac. D60 changes every 30 arcminutes. A small longitude change can therefore produce a different varga placement even when D1 remains unchanged. Birth-time error can be just as important as ayanamsa, especially for the Ascendant and house cusps.

Should a transit website display its ayanamsa?

Yes. It should also display the timestamp, timezone, geocentric or topocentric setting, node type, and retrograde convention. A date without those settings looks precise while remaining difficult to reproduce.

Will a future astronomical model force all ayanamsas to agree?

No. Better astronomy can improve how each chosen anchor is calculated, but it cannot make Spica, Delta Cancri, Zeta Piscium, Lambda Scorpii, and the Galactic Center become the same reference. The remaining disagreement is about the astrological convention itself.

What should I do if my Moon or Ascendant changes between Lahiri and True Chitra?

Treat the chart as boundary-sensitive. Verify the birth record, time standard, location, and software settings. Then compare predictions prospectively under both systems. Do not rectify the birth time merely to preserve a preferred sign. A one-minute ayanamsa difference can be meaningful at a boundary without making the entire chart unreliable.

A CLEAR RECOMMENDATION

16. The conclusion: which ayanamsa makes the most sense in 2026?

If I must choose one ayanamsa for a general Vedic astrology practice in 2026, I choose Lahiri IAE 1985.

The reason is not that astronomy has crowned Lahiri or that every alternative has failed. The reason is that an ayanamsa is part astronomical definition and part astrological convention. Standard Lahiri has the strongest combination of reproducibility, Indian calendar continuity, software support, published interpretive literature, teacher-to-student continuity, and compatibility with the charts most Vedic astrologers are already discussing.

True Chitra is the strongest precision companion. If your principle is literally that Spica must remain at 0 degrees Libra, True Chitra implements that principle more faithfully than an epoch-based approximation. In 2026 it differs from standard Lahiri by only about 1 minute 11 seconds. That makes it ideal for a sensitivity check, especially for the Moon, Ascendant, KP-style fine divisions, and high Vargas near boundaries.

Pushya, Raman, True Mula, True Revati, Dhruva, and galactic systems are meaningful when used inside the methods and philosophical frameworks that motivated them. They should remain available as clearly named research options. They should not silently replace Lahiri in a general Vedic app, because the differences can move nakshatras, padas, dashas, and transit dates far enough to break comparison with mainstream practice.

Final ruling for 2026
Default: Lahiri IAE 1985.
Precision check: True Chitra.
KP: the exact named KP variant.
Alternative schools: use the system required by the method and disclose it.
Universal predictive winner: not yet established.

Ayanamsa should make a chart more transparent, not more mystical. Name the grid, keep the calculation stable, flag the boundaries, and judge techniques on predictions recorded before the outcome. That approach respects both tradition and honest observation.

FOR TECHNICAL COMPARISON

Appendix A. Extended 2026 values

All values are mean ayanamsa at 2026-08-27 18:39:49 UTC, rounded to 0.01 arcsecond. The delta is measured from Lahiri. Letter d means degrees, m means arcminutes, and s means arcseconds.

System	Mean ayanamsa	Delta from Lahiri
Fagan-Bradley	25d 06m 45.52s	+52m 59.5s
Lahiri IAE 1985	24d 13m 45.97s	0m 00.0s
Lahiri ICRC pre-1985	24d 13m 44.88s	-0m 01.1s
Lahiri 1940	24d 12m 52.80s	-0m 53.2s
Lahiri VP285	24d 14m 08.97s	+0m 23.0s
True Chitra	24d 12m 34.87s	-1m 11.1s
KP table	24d 07m 57.31s	-5m 48.7s
KP VP291	24d 09m 09.75s	-4m 36.2s
Raman	22d 46m 59.29s	-1d 26m 46.7s
True Pushya	23d 05m 22.57s	-1d 08m 23.4s
True Revati	20d 25m 17.03s	-3d 48m 28.9s
Surya Siddhanta equinox	21d 16m 02.65s	-2d 57m 43.3s
Aryabhata equinox	21d 16m 02.65s	-2d 57m 43.3s
Yukteswar	22d 51m 04.13s	-1d 22m 41.8s
GC at 0 Sagittarius	27d 13m 34.19s	+2d 59m 48.2s
Dhruva GC Middle Mula	20d 26m 10.14s	-3d 47m 35.8s
True Mula	24d 57m 35.83s	+43m 49.9s
Gil Brand	22d 50m 57.20s	-1d 22m 48.8s
Mardyks	30d 23m 24.50s	+6d 09m 38.5s
Fiorenza	25d 22m 20.51s	+1d 08m 34.5s
Ardra Galactic Plane	23d 47m 01.52s	-26m 44.5s
Usha-Shashi	20d 25m 47.59s	-3d 47m 58.4s
Sassanian	20d 21m 55.09s	-3d 51m 50.9s
Aldebaran at 15 Taurus	25d 07m 52.56s	+54m 06.6s
Babylonian Huber	25d 06m 21.53s	+52m 35.6s
Babylonian Britton	24d 59m 17.14s	+45m 31.2s

PLAIN-LANGUAGE DEFINITIONS

Appendix B. Glossary

Term	Meaning
Ayanamsa	The angular offset between the tropical zodiac and a selected sidereal zodiac.
Chitrapaksha	The family of sidereal systems oriented to Chitra, commonly identified with Spica.

Term	Meaning
Ecliptic	The apparent annual path of the Sun and the reference plane used for zodiac longitude.
Ephemeris	A table or computation of celestial positions at specified times.
Fiducial	The reference object or rule used to define the zodiac's zero point.
Geocentric	Calculated as seen from Earth's center. Most standard transit tables use this frame.
Mean ayanamsa	An offset relative to the mean equinox, without short-period nutation.
Nakshatra	One of 27 equal lunar mansions, each spanning 13 degrees 20 minutes.
Nutation	A shorter-period oscillation added to the longer precessional motion.
Pada	One quarter of a nakshatra, spanning 3 degrees 20 minutes and mapping to Navamsha.
Precession	The long-term change in the orientation of Earth's equator and related coordinate origins.
Proper motion	A star's gradual change of position relative to the solar system's reference frame.
Sidereal zodiac	A twelve-sign zodiac whose origin is tied to a stellar, historical, or geometric rule.
Tropical zodiac	A twelve-sign zodiac beginning at the moving March equinox.
True Chitra	A dynamic ayanamsa that keeps Spica exactly at 0 degrees Libra by recomputing the star.
Varga	A divisional chart created by mapping fractions of each sign into another zodiac.

About the author

Agastya Trivedi is a Vedic astrologer focused on making classical ideas understandable without hiding the assumptions inside modern calculations. His approach combines traditional technique, careful software practice, and an insistence on naming uncertainty where it genuinely exists.

This guide is educational. It presents astrological traditions and astronomical definitions, not scientific proof that astrology can predict individual outcomes.